

5. VERY LOW- TO LOW-GRADE METAMORPHIC ROCKS

Recommendations by the IUGS Subcommittee on the Systematics of Metamorphic Rocks.
Web version of 01/02/07

Péter Árkai¹, Francesco P. Sassi² and Jacqueline Desmons³

¹ P. Árkai, Institute for Geochemical Research, Hungarian Academy of Sciences, Budapest, Hungary

² Department of Mineralogy and Petrography, University of Padova, Italy

³ CNRS, Nancy, France

Introduction

A Study Group was set up in 1987 under the leadership of P. Árkai to study the nomenclature and systematics of rocks and processes related to the so-called very low- to low-grade metamorphism, that is, from the diagenetic field to the greenschist facies or epizone. The Study Group consisted of 30 scientists from 5 continents who were specialists in this area and who agreed to cooperate. This paper presents the definitions that were agreed, by repeated iterations, considering on one hand the various opinions within the Study Group (largely gathered through questionnaires), and by discussing these results within the SCMR, on the other. Being the result of multiple discussions and compromises, the definitions presented here are recommended for international use.

Very low- and low-grade metamorphism

The SCMR agreed that for the purposes of its discussions and definitions the grade of metamorphism would be taken as equivalent to the temperature of metamorphism. The whole temperature/grade range of metamorphism has been divided into five parts, namely: very low, low, medium, high and very high (Smulikowski et al., this vol.) The present paper deals with the nomenclature of the very low- and low-grade rocks. It also covers the transition from diagenesis to metamorphism and from very low grade to low grade (the greenschist facies or epizone).

From diagenesis to metamorphism

Diagenesis

According to the SCMR (Smulikowski et al., this vol.), metamorphism is: ‘*a process involving changes in the mineral content/composition and/or microstructure of a rock, dominantly in the solid state. This process is mainly due to an adjustment of the rock to physical conditions that differ from those under which the rock originally formed and that also differ from the physical conditions normally occurring at the surface of the Earth and in the zone of diagenesis. The process may coexist with partial melting and may also involve changes in the bulk chemical composition of the rock*’. Consistent with this definition, the term diagenesis covers the lowest temperature part of the changes in the outer part of the Earth’s crust, excluding weathering.

Diagenesis (*sensu lato*): *All the chemical, mineralogical, physical and biological changes undergone by a sediment after its initial deposition, and during and after its lithification, exclusive of superficial alteration (weathering) and metamorphism.* The changes involved in diagenesis are the result of such processes as compaction, cementation, reworking, authigenesis, replacement, crystallisation, leaching, hydration, dehydration, bacterial action, and formation of concretions. These processes occur under conditions of pressure and temperature that are normal at the Earth’s surface and in the outer part of the Earth’s crust.

Diagenesis (*sensu lato*) may be subdivided into:

- shallow diagenesis (=diagenesis *sensu stricto*): *the chemical, mineralogical, physical and biological changes that take place in a sediment under physical conditions that do not differ significantly from those under which the sediment originated. It is characterised by the absence of alteration of detrital minerals.*
- deep diagenesis: *changes that are characterised by clay mineral reactions (such as the transformation of smectite to illite, kaolinite to dickite, etc., and the increase of the proportion of illite layers in interstratified clay minerals).*

Instead of shallow and deep diagenesis the adjectives ‘early’ and ‘late’ are also used in the literature. However, in order to avoid the time connotation implied by these adjectives, the SCMR prefers the terms ‘shallow’ and ‘deep’.

Deep diagenesis is the term recommended by the SCMR, as the equivalent to the three terms epigenesis, katagenesis and catagenesis of Russian authors, and middle + late or deep burial stage diagenesis of Müller (1967) and Dunoyer de Segonzac (1970).

Very low-grade metamorphism: the transition zone

The transition zone between diagenesis (*sensu lato*) and metamorphism, effectively the field of very low-grade metamorphism, is characterised by the gradual change of various features, which affect the rocks through this zone up to their partial or complete alteration into metamorphic rocks.

However, in the context of rock nomenclature, the main problem is that the most critical changes, marking this transition, are not visible with the naked eye because they commonly occur only on the microscopic or submicroscopic scale. Thus, the related nomenclature

requires an important exception to the SCMR principle (Schmid et al., this vol.), which states that, wherever possible, definitions are based on features that can be seen with the naked eye.

Indeed, on a mesoscopic scale these transitional rocks commonly display characteristics identical with, or very similar to, those of their non-metamorphic equivalents. Therefore, these very low-grade rocks can, in some cases, only be recognised by means of microscopic investigations and in most cases using other instrumental techniques. These include measurements of the illite Kübler index (formerly called 'crystallinity') and coalified organic matter order-disorder by means of X-ray powder diffraction, measurements of vitrinite reflectance by means of optical microscopy, fluid inclusion thermobarometry, etc.

Different criteria have been used for the characterisation and subdivision of the transitional field between diagenesis and low-grade metamorphism in various rock types. Consequently, various systems and nomenclatures have been established (e.g. specific mineral associations, illite Kübler index zones, microstructural zones, coal rank scales, etc.). See Figure 5.1 for a schematic comparison of these systems. Details and comprehensive interpretations can be found in the textbooks edited by Frey (1987a) and Frey & Robinson (1999), including the studies of Frey (1987b), Liou et al. (1987), Teichmüller (1987), Mullis (1987), Merriman & Peacor (1999), Merriman & Frey (1999), Robinson & Bevins (1999) and Alt (1999).

The temperature (and pressure) boundaries related to the sequence of stages or zones of each nomenclature scheme are not defined exactly, and in most cases they do not coincide with those of the other schemes. The correlation between the nomenclature schemes and absolute temperature is full of uncertainties, mostly because of the greatly differing nature of the disequilibrium processes considered, the transitional open- semi-closed systems, and in consequence, the great variations in the chemical effects of the fluids present.

Because the transition between diagenesis and metamorphism is gradual, any boundary between these two fields can only be arbitrarily defined. In practice, it is not possible to establish an isothermal boundary that can be applied to all, or even to the majority, of the rock types. Furthermore, rocks having different bulk composition, different grain-size, different amount of strain, different porosity and permeability, etc., may react at relatively higher or lower temperatures, so that, at a given depth or temperature a given rock may react whilst others may remain unchanged. Thus, the temperature of the beginning of metamorphism may vary considerably between rock types.

The problem of mixed sequences

The problem of different rock types, as discussed above, raises a related and important question. If the very-low grade character of a rock in a lithologically mixed sequence can be determined (e.g. as the prehnite-pumpellyite facies (see below) in basic volcanic rocks): can that classification be extrapolated to other rock types in the sequence even though they do not display any measurable alteration in hand specimen (and, commonly, not even at the microscopic scale)? Alternatively, should different grade terms be used for the different rock types?

To put the question another way. Within the same rock pile, depending on the chemical, mineralogical and physical properties of the different rocks, some (e.g. carbonate rocks) may show hardly any signs of alteration, whereas others that underwent the same T-P_l-P_f¹

¹ P_l – lithostatic pressure; P_f – fluid pressure
Paper_5.doc

conditions, are clearly recrystallised, and therefore must be classified as metamorphic rocks, and thus receive metamorphic rock names. In these cases, should the metamorphic name/grade be extrapolated to the unaltered rocks, or should they be given the most appropriate, non-metamorphic name, disregarding that of their surroundings?

In response to this very difficult question the SCMR recommends the following:

- *those rocks which display at least one observable or measurable sign of very low-grade metamorphism (i.e. diagnostic minerals or mineral assemblages, appropriate values of illite Kübler index, appropriate values of vitrinite reflectance, etc.) should be given metamorphic rock names;*

- *those rocks which do not show any sign of very low-grade metamorphism should be given the appropriate non-metamorphic rock name, regardless of their close spatial and genetic relationship to the 'metamorphic' rocks.*

This implies that metamorphic rock names may alternate with sedimentary or magmatic ones in a geologic profile or map. This might give rise to cartographic boundaries that would require specific explanations.

The various nomenclature systems of very low- and low-grade metamorphism

The possibilities for discrimination between diagenesis, very low- and low-grade metamorphism are schematically shown in Figure 5.1.

Rocks with diagnostic minerals and mineral assemblages (mainly basic rocks)

a) Zeolite and subgreenschist facies

In rocks of basic to intermediate compositions, the occurrence of zeolites is attributed to alterations at low temperatures in the presence of CO₂-poor or -absent aqueous fluids. The SCMR defines the zeolite facies as follows:

Zeolite facies: *a facies (in the sense of Eskola, 1920) that embraces all mineral assemblages that include various zeolites plus quartz, irrespective of the mode of origin, whether metamorphic, (including hydrothermal) or diagenetic (Coombs et al., 1959; Coombs, 1971; Boles & Coombs, 1977).*

In silicate rocks (see Bucher & Frey, 1994), the first appearance of one or some of the following minerals: Fe-Mg-carpholite, glaucophane, lawsonite, laumontite, paragonite, prehnite, pumpellyite or stilpnomelane, is commonly regarded as the beginning of metamorphism (although the status of laumontite is questionable²).

² Coombs et al. (1959) and Boles & Coombs (1977) argue that the zeolite facies should be regarded as a *mineral facies*, irrespective of the origin of the mineral assemblage. They regarded it as unrealistic to arbitrarily interpret laumontite as metamorphic and heulandite or analcime (plus quartz) as diagenetic.

The SCMR recommends that only a few metamorphic facies should be used as a general rule (Smulikowski et al., this vol.), but leaves open the possibility to use other facies or subfacies if required to describe a specific region, provided that these additional facies or subfacies are clearly defined. Although the very low- and low-grade P-T field covers quite a small part of the whole metamorphic field, various authors have suggested that it should be subdivided into several facies or subfacies to cover the situation with particular lithologies. They also suggest different methods for defining the additional facies or subfacies.

The following are a few of these suggestions and the recommended SCMR definitions. As grade is taken as equivalent to temperature (Smulikowski et al., this vol.) it follows that metamorphic rocks formed at temperatures lower than the low-temperature limit of the greenschist facies belong to the very low-grade field. The high P/T and high-P part of this field includes a small part of the glaucophane-schist (blueschist) facies (Smulikowski et al., this vol.).

The part of the field of very low-grade metamorphism characterised by pressures lower than those of the glaucophane-schist facies has been called the **subgreenschist facies** (e.g. Bucher & Frey, 1994; Merriman & Frey, 1999). The subgreenschist facies embraces various metabasite facies/subfacies (Fig. 5.1) characterised by the diagnostic mineral assemblages of prehnite + pumpellyite, prehnite + actinolite, pumpellyite + actinolite, and also, according to certain authors, laumontite.

The SCMR defines these facies/subfacies as follows.

*The **prehnite-pumpellyite facies** is characterised in metasediments and metavolcanic rocks of appropriate composition by the presence of prehnite and/or pumpellyite in the absence of zeolites, lawsonite, or jadeite. Quartz-albite-chlorite-prehnite and/or pumpellyite may coexist stably (Coombs, 1960).*

*The **pumpellyite-actinolite facies** is characterised by the mineral association of pumpellyite-actinolite-quartz (\pm chlorite, albite and epidote) and by the lack of prehnite (Hashimoto, 1966).*

*The **prehnite-actinolite facies** is characterised by the mineral association of prehnite-actinolite-epidote (\pm chlorite, albite, quartz and titanite) and by the absence of pumpellyite in rocks of appropriate bulk composition (mostly metabasic rocks and their clastic derivatives) (Liou et al., 1985).*

b) The transition to the greenschist facies

In the case of rocks of suitable composition (mostly of basic to intermediate composition), the first appearance of the diagnostic mineral assemblage actinolite + epidote + chlorite + albite + quartz in the absence of pumpellyite and/or prehnite indicates the onset of low-grade metamorphism, that is, the transition from very low- to low-grade metamorphism, or from subgreenschist to greenschist facies.

The first appearance of the lawsonite + chlorite + albite association or of sodic amphibole lies within the very low-grade field, indicating high P/T.

Although very fine-grained chloritoid may appear in very low-grade (anchizonal) metapelites (commonly hard to detect by optical microscopy), chloritoid typically occurs in low-grade (greenschist facies) rocks.

Rocks devoid of diagnostic minerals and mineral assemblages

In many common rocks, such as 'normal' marine pelites, carbonate rocks, etc., no diagnostic minerals and mineral assemblages form in the very low-grade field. In these rocks, the transitions from non-metamorphic to very low-grade and from very low-grade to low-grade metamorphic domains take place through the diagenetic zone, the anchizone and the epizone (Fig. 5.1), each zone being characterised by specific values of the illite Kübler index (KI), which is measured on the $<2\ \mu\text{m}$ fraction of clay-rich clastic rocks following the recommendations on sample preparation, X-ray diffraction settings and inter-laboratory standardisation summarised by Kisch (1991). For comparing the various sample preparation techniques, the procedure and standards suggested by Warr & Rice (1994) are useful. Concerning the proper nomenclature of indices (formerly called 'crystallinity') expressing the reaction progress of illite-muscovite and chlorite in diagenetic and low-temperature metamorphic conditions, the authors refer to the recommendations of Guggenheim et al. (2002).

Diagenetic zone [or more precisely, *diagenetic illite Kübler index zone*]: zone characterised by illite Kübler index (KI) values greater than $0.42\ \Delta^{\circ}2\theta\ \text{CuK}\alpha$ (after Kübler, 1967, 1968, 1984).

Anchizone: transitional zone between the diagenetic zone and the epizone characterised by illite Kübler index (KI) mean values between 0.42 and $0.25\ \Delta^{\circ}2\theta\ \text{CuK}\alpha$ (after Kübler, 1967, 1968, 1984). Metamorphism in this zone is consistently called **anchimetamorphism**, which roughly corresponds to very low-grade metamorphism. Note: the term 'anchimetamorphism' was originally introduced by Harsowitz (1927) to indicate changes in mineral content of rocks under pressure and temperature conditions prevailing between the Earth's surface and the zone of metamorphism.

Epizone: zone of low-grade metamorphic rocks characterised by illite Kübler index (KI) mean values less than $0.25\ \Delta^{\circ}2\theta\ \text{CuK}\alpha$ (after Kübler, 1967, 1968, 1984). Note: the term epizone was originally proposed by Grubenmann (1904) to indicate shallow depth of metamorphism. At present, however, this term is mainly used in the context of illite Kübler index investigations.

In addition to the illite Kübler-index, some other characteristics such as vitrinite reflectance (Fig. 5.1), chlorite Árkai index (see Guggenheim et al., 2002), Conodont Colour Alteration Index (CAI), etc. can also be used for determining diagenetic and metamorphic zones. Note, however, that only approximate correlations can be established between these parameters. Examples and explanations of common deviations from the generalised scheme are given by Kisch (1987) and Merriman & Frey (1999).

Protolith names, definitions of specific rock names

Very few specific rock terms exist in the realm of very low-grade metamorphism. This can be explained by the fact that, with the exception of the structure of slates, mesoscopic features characteristic of this metamorphic grade do not exist.

As the original (sedimentary or magmatic) features of the protoliths can usually be easily recognised in this realm, the general use of the sedimentary or magmatic name of the protolith prefixed with 'meta' or 'meta-' is highly recommended (e.g. meta-andesite, metasandstone), excluding the very few cases when *specific rock terms* are available (e.g. slate). The SCMR recommends the use of the *prefix meta or meta-* in combination with a protolith-based name only when the protolith is easily identifiable or obvious. It must never be used for former metamorphic rocks (for example, 'meta-phyllite' is not an acceptable term) (Schmid et al., this vol).

The *specific rock terms* are defined below.

Slate: *An ultrafine- or very fine-grained rock displaying slaty cleavage.* Slate is usually of very low metamorphic grade, although it may also occur in low-grade conditions. The definition of *slaty cleavage* is given by Brodie et al. (this vol.) as; '*A type of continuous cleavage in which the individual grains are too small to be seen by the unaided eye*'. The definition of *continuous cleavage* is given by Brodie et al. (this vol.) as '*A type of cleavage characterised by the preferred orientation of all inequant mineral constituents of a rock, and in which the cleavage planes are developed at the grain size scale*'.

Phyllite: *A fine- to medium-grained rock characterised by a lustrous sheen and a well-developed schistosity resulting from the parallel arrangement of phyllosilicates. Phyllite is usually of low metamorphic grade.*

Palaeovolcanite rock terms: *the SCMR recommends following the definitions for volcanic rock terms given by Le Maitre et al. (1989, 2002) and that the recommendations given in the present paper are used for rocks which underwent any kind of very low-grade metamorphic alteration.*

Spilite: *An altered basic to intermediate, volcanic or subvolcanic rock in which the feldspar is partially or completely composed of albite and is typically accompanied by chlorite, calcite, quartz, epidote, prehnite, or other low-temperature hydrous crystallisation products.* Preservation of eruptive (volcanic and subvolcanic) features is an important characteristic of spilites. The term spilite may be classified as metabasalt or meta-andesite, as appropriate, regardless of its origin.

The 'spilite problem', debated in the geological literature for many decades, is of special interest for researchers working in the field of very low- and low-grade metamorphism. This is the reason why the term 'spilite' has been considered by the SCMR, despite the fact that the Subcommittee on the Systematics of Igneous Rocks include this term in their glossary³ (Le Maitre et al., 1989, 2002). The SCMR recommends that the definition given by Le Maitre et al. should be expanded to include rocks of intermediate composition, because many spilites have an andesitic rather than a basic composition.

Greenschist: *Schist whose greenish colour is due to the presence of minerals such as actinolite, chlorite and epidote.* More precise terms should be used whenever possible (e.g. epidote bearing actinolite-chlorite schist).

Greenstone: *A granofels whose greenish colour is due to the presence of minerals such as actinolite, chlorite and epidote.* More precise terms should be used whenever possible (e.g. chlorite-epidote granofels).

² Note: the term 'spilite' is not a root or recommended name in the IUGS classification of Igneous Rocks, although it is included in the glossary (Le Maitre, 2002)
Paper_5.doc

Terms not recommended for general use

In addition to the terms listed above, the Study Group also discussed some other terms, even though they are not recommended for international use. They are listed below, with the corresponding explanations.

Brownstone facies: *a low-temperature mineral facies encompassing ocean-floor weathering and low-temperature hydrothermal alteration of the ocean floor.* The most widespread secondary minerals present under oxidising conditions include a K-rich dioctahedral iron illite resembling celadonite. This replaces olivine, occupies vesicles, replaces interstitial glass, and eventually, augite. Under reducing conditions its place is taken by saponite, and pyrite is also characteristic. Plagioclase may be replaced by clay minerals or potassium feldspar. Glassy rinds of basaltic pillow lava alter to palagonite in association with phillipsite and other low-temperature zeolites and calcite.

Catagenesis=katagenesis: *terms used, especially by Russian authors (see above), to indicate changes occurring in (an already lithified) sedimentary rock buried under a distinct covering layer, characterised by P-T conditions that are significantly different from those of both deposition and metamorphism.* This term is equivalent to 'epigenesis' or to 'middle and deep diagenesis'. It is subdivided into early catagenesis (= middle diagenesis = shallow epigenesis) and late catagenesis (= deep diagenesis = deep epigenesis).

Cryptic metamorphism: *a term for metamorphism that can be detected only by special study, for example, vitrinite reflectance, illite Kübler index, etc. and not by ordinary hand specimen or microscopic study.*

Crystalline limestone, dolomite, etc.: *carbonate rock (carbonate mineral content > 90 %) with a completely recrystallised matrix.* Foliation may be present.

Epigenesis: *a term used, especially by Russian authors, to indicate changes, transformations or processes, occurring at low temperatures and pressures that affect sedimentary rocks after their compaction, excluding superficial alteration (weathering) and metamorphism: (= catagenesis = middle and deep diagenesis).* Subdivided into early and deep or late epigenesis.

Incipient regional metamorphism: *a general term for the stages of mineral modifications as characterised by the appearance of the attributes of the anchizone (anchimetamorphism).* It is approximately equivalent to the early metagenesis or with the greater part of Winkler's (1974) 'very low-grade metamorphism', including the higher-T part of the 'pumpellyite-prehnite-quartz facies', and probably, all of the 'lawsonite-albite schist' and 'glaucophane-lawsonite schist facies' in the sense of Winkler (1974).

Metagenesis: *a term used, especially by Russian authors (see above), to indicate a more advanced stage of post-diagenetic alteration than epigenesis or catagenesis.* It is subdivided into 'early metagenesis' (which roughly corresponds to the anchizone) and 'late or deep metagenesis' (which is more or less equivalent to the epizone or the chlorite zone of the greenschist facies).

ACKNOWLEDGEMENTS

The valuable help of all colleagues in the Study Group on “Very Low- and Low-Grade Rocks” is gratefully acknowledged. The following Study Group members have substantially contributed to the elaboration of the first, preliminary draft of the present paper: K. Balogh, S. Banno, R.J. Bevins, T. Blenkinsop, C. Brime, K. Brodie, D.S. Coombs, L. Cortesogno, N.J. Fortey, M. Frey, E.D. Ghent, M. Hashimoto, A. Iijima, H.J. Kisch, B. Kübler, D. Laduron, J.G. Liou, F. Lippmann, N.V. Logvinenko, J. Martini, S. Morad, R. Offler, L.P. Plyusnina, D. Robinson, B. Roberts, P. Schiffman, V. Suchy, M. Vuagnat and C.E. Weaver. Valuable comments and corrections on the Web version 30.10.2002 given by S. Banno, T. Blenkinsop, C. Brime, D.S. Coombs, H. Day, D.D. Eberl, E. Essene, N. Fortey, H.J. Kisch, L. Leoni, J.G. Liou, R.J. Merriman, F. Nieto, R. Offler, B. Roberts and P. Schiffman are gratefully acknowledged. Thanks are due to the IUGS for partially supporting the annual working meetings of the SCMR.

The work of P.Á. has also been supported by the Hungarian National Science Foundation (OTKA, Budapest), Projects Nos. T007211/1993-1996, T022773/1997-2000 and T035050/2001-2004. F.P.S. acknowledges support by Italian CNR (Istituto di Geoscienze e Georisorse) and MIUR.

REFERENCES

- Alt J.C. 1999. Very low-grade hydrothermal metamorphism of basic igneous rocks. In: Low-Grade Metamorphism, M. Frey & D. Robinson, (Eds.), Blackwell Science, Oxford, 169-201.
- Bates R. L. and Jackson J. A. 1987. Glossary of Geology (3rd edition). American Geological Institute. Alexandria, Virginia.
- Boles J. R. and Coombs D. S. 1977. Zeolite facies alteration of sandstones in the Southland Syncline, New Zealand. *Amer. J. Sci.* 277, 982-1012.
- Brodie K., Fettes, D.J. Harte, B. and SCHMID, R. 2007, A systematic nomenclature for metamorphic rocks: 3. Structural terms including fault rocks. Recommendations by the IUGS Subcommittee on the Systematics of Metamorphic Rocks. SCMR website (www.bgs.ac.uk/SCMR).
- Bucher K. and Frey M. 1994. Petrogenesis of metamorphic rocks. 6th edition. Springer-Verlag, Berlin, 318 pp.
- Coombs D. S. 1960. Lower grade metamorphic facies in New Zealand. 21st Int. Geol. Congr., Copenhagen, 1960, 13, 339-351.
- Coombs D. S. 1971. Present status of the zeolite facies. In: Molecular sieve zeolites, 1, *Advances in Chemistry Ser.* 101. Am. Chem. Soc. 317-327.
- Coombs D. S., Ellis A. J. Fyfe W. S. and Taylor A. M. 1959. The zeolite facies, with comments on the interpretation of hydrothermal syntheses. *Geochem. Cosmochem. Acta* 17, 53-107.
- Dunoyer de Segonzac G. 1970. The transformation of clay minerals during diagenesis and low-grade metamorphism: a review. *Sedimentology* 15, 281-346.
- Eskola P. 1920. The mineral facies of rocks. *Norsk. Geol. Tidsskr.* 6, 143-194.
- Frey M. ed. 1987a. Low temperature metamorphism. Blackie, Glasgow and London, p.351.
- Frey M. 1987b. Very low-grade metamorphism of clastic sedimentary rocks. In: Low Temperature Metamorphism, M. Frey (Ed.), Blackie, Glasgow and London, 9-58.
- Frey M. and Robinson D. (Eds.). 1999. Low-grade metamorphism. Blackwell Science, Oxford, p.328.
- Grubenmann U. 1904. Die kristallinen Schiefer. Bornträger, Berlin.
- Guggenheim S., Bain D. C., Bergaya F., Brigatti M. F., Drits V. A., Eberl D. D., Formoso M. L. L., Galán E., Merriman R. J., Peacor D. R., Dtanjek H., Watanabe T. 2002. Report of the Association internationale pour l'étude des argiles (AIPEA) Nomenclature Committee for 2001: Order, disorder and crystallinity in phyllosilicates and the use of the 'crystallinity index'. *Clays and Clay Minerals*, 50, 406-409.
- Harrasowitz H. 1927. Anchimetamorphose, das Gebiet zwischen Oberflächen- und Tiefenumwandlung der Erdrinde. *Ber. Oberhess. Ges. Nat. Heilk. Giessen, Naturwiss. Abt.*, 12, 9-15.
- Hashimoto M. 1966. On the prehnite-pumpellyite metagraywacke facies. *J. Geol. Soc. Japan*, 72, 253-265.

- Kisch H. J. 1987. Correlation between indicators of very low-grade metamorphism. in: "Low Temperature Metamorphism", M. Frey (Ed.), Blackie, Glasgow and London, 227-300.
- Kisch H. J. 1991. Illite crystallinity: recommendations on sample preparation, X-ray diffraction settings, and interlaboratory samples. *J. metamorphic Geol.* 9, 665-670.
- Kisch H.J., Árkai P., Covadonga B. 2004. Calibration of illite Kübler index (illite "crystallinity"). *Schweizerische Mineralogische und Petrographische Mitteilungen*, 84, 323-331.
- Kübler B. 1967. La cristallinité de l'illite et les zones tout a fait superieures du métamorphisme. In: *Étages tectoniques, Colloque de Neuchâtel 1966, A La Baconniere, Neuchâtel* 105-121.
- Kübler B. 1968. Evaluation quantitative du métamorphisme par la cristallinité d'illite. *Bull. Centre Rech. Pau - S.N.P.A.* 2, 385-397.
- Kübler B. 1984. Les indicateurs des transformations physiques et chimiques dans la diagenèse, température et calorimétrie. in: "Thermobarométrie et barométrie géologiques", M. Lagache ed., Soc. Franc. Minéral. Cristallogr., Paris, 489-596.
- Le Maitre R. W. (Ed.) et al., 1989. A classification of igneous rocks and glossary of terms. Blackwell Sci. Publ., 193p.
- Le Maitre R. W. (Ed.) et al., 2002. A classification of igneous rocks and glossary of terms. Cambridge University Press, 236p.
- Liou J.G., Maruyama S. and Cho M. 1985. Phase equilibria and mineral parageneses of metabasites in low-grade metamorphism. *Miner. Mag.*, 49, 321-333.
- Liou J.G., Maruyama S. and Cho M. 1987. Very low-grade metamorphism of volcanic and volcanoclastic rocks – mineral assemblages and mineral facies. in: *Low Temperature Metamorphism*, M. Frey, ed., Blackie, Glasgow and London, 59-113.
- Merriman R.J. and Frey M. 1999. Patterns of very low-grade metamorphism in metapelitic rocks. In: *Low-Grade Metamorphism*, M. Frey & D. Robinson (Eds.), Blackwell Science, Oxford, 61-107.
- Merriman R. J. and Peacor D. R. 1999. Very low-grade metapelites: mineralogy, microfabrics and measuring reaction progress. In: *Low-Grade Metamorphism*, M. Frey & D. Robinson (Eds.), Blackwell Science, Oxford, 10-60.
- Mullis J., 1987. Fluid inclusion studies during very low-grade metamorphism. in: „Low Temperature Metamorphism”, M. Frey, ed., Blackie, Glasgow and London, 162-199.
- Müller G., 1967. Diagenesis in argillaceous sediments. In: *Diagenesis in Sediments*, Larsen G. and Chilingar G. V. eds., Elsevier, Amsterdam, pp. 127-177.
- Robinson D. and Bevins R.E., 1999. Patterns of regional low-grade metamorphism in metabasites. In: *Low-Grade Metamorphism*, M. Frey & D. Robinson (Eds.), Blackwell Science, Oxford, 143-168.
- Rosen O. and Callegari E., (2003.): Towards a unified nomenclature in metamorphic petrology: 7. Carbonate and related rocks. A proposal on behalf of the IUGS Subcommittee on the Systematics of Metamorphic Rocks. SCMR website (www.bgs.ac.uk/SCMR).
- Schmid R., Fettes D., Harte B., Davis E., Desmons J., Meyer-Marsilius H-J. and Siivola J. 2004, A systematic nomenclature for metamorphic rocks: 1. How to name a metamorphic rock. Recommendations by the IUGS Subcommittee on the Systematics of Metamorphic Rocks. SCMR website (www.bgs.ac.uk/SCMR).
- Smulikowski W., Desmons J., Harte B., Sassi F.P. and Schmid R. 2007, A systematic nomenclature for metamorphic rocks: 2. Types, grade and facies of metamorphism. Recommendations by the IUGS Subcommittee on the Systematics of Metamorphic Rocks. SCMR website (www.bgs.ac.uk/SCMR).
- Teichmüller M. 1987. Organic material and very low-grade metamorphism. In: *Low Temperature Metamorphism*, M. Frey (Ed.), Blackie, Glasgow and London, 114-161.
- Warr L.N., Rice A.H.N. 1994. Interlaboratory standardization and calibration of clay mineral crystallinity and crystallite size data. *Journal of Metamorphic Geology*, 12, 141-152.
- Wilson, G. 1961. The tectonic significance of small scale structures, and their importance to the geologist in the field. *Annales de la Soc. Géol. de la Belgique*, 84, pp423-548.
- Winkler H.G.F. 1974. *Petrogenesis of Metamorphic Rocks*. (3rd edition). Springer-Verlag, New York

Captions:

Fig. 5.1: Comparison of mineral facies, illite Kübler index (KI) 'crystallinity' zones and coal rank in the diagenetic, very low- and low-grade metamorphic realms. Zigzag lines represent uncertainties of correlation. Rough scheme simplified after Frey (1987), Kisch (1987) and Merriman & Frey (1999).

metamorphic grade	mineral facies	illite K _{bler} -index (KI) zone	coal rank, R _{random} % vitrinite reflectance
diagenesis	zeolite facies (incl. laumontite zone)	diagenetic zone (KI > 0.42Δ20)	bituminous coal
very low-grade	prehnite-pumpellyite facies	anchizone (KI = 0.42 - 0.25Δ20)	2.0
	subgreenschist facies prehnite-actinolite facies pumpellyite-actinolite facies <i>increasing pressure</i>		2.5
low-grade	greenschist facies	epizone (KI < 0.25Δ20)	4.0 - 5.0

glaucophane schist facies

Fig 5.1